

Cramér Type Moderate Deviation for the Maximum of the Periodogram with Application to Simultaneous Tests in Gene Expression Time Series

Weidong Liu¹ and Qi-Man Shao²

Hong Kong University of Science and Technology

Abstract. In this paper, Cramér type moderate deviations for the maximum of the periodogram and its studentized version are derived. The results are then applied to a simultaneous testing problem in gene expression time series. It is shown that the level of the simultaneous tests is accurate provided that the number of genes G and the sample size n satisfy $G = \exp(o(n^{1/3}))$.

Keywords: moderate deviation, periodogram, simultaneous tests.

AMS 2000 subject classification: Primary 60F05.

1 Introduction

Let $X_1, X_2, \dots$ be a sequence of random variables. Define the periodogram ordinates for $\{X_n\}$ at the standard frequencies $\omega_j = 2\pi j/n$ by

$$I_n(\omega_j) = \frac{1}{n} \left| \sum_{k=1}^n X_k e^{ik\omega_j} \right|^2,$$

where $1 \leq j \leq q$ and $q = [(n-1)/2]$.

The periodogram is a fundamental tool in spectral analysis and is often used to detect periodic patterns in various real applications, such as the analysis of gene expression data and the study of earthquake. Theoretical properties of the periodogram have been extensively studied. An, Chen and Hannan (1983) obtained the logarithm law for the maximum of the periodogram; Davis and Mikosch (1999), Mikosch, Resnick and Samorodnitsky (2000), Lin and Liu (2009a) obtained the asymptotic distribution for the maximum of the periodogram under the i.i.d. and linear process cases, the heavy-tailed case and nonlinear time series case respectively; Fay and Soulier (2001) obtained central limit theorems for functionals of the periodogram; Shao and Wu (2007) obtained asymptotic distributions for the periodogram and empirical distribution function of the periodogram for a wide class of nonlinear processes.

¹Email: liuweidong99@gmail.com

²Email: maqms hao@ust.hk. Research partially supported by Hong Kong RGC CERG 220608

When $\{X_n\}$ are independent and identically distributed (i.i.d.) random variables with $\text{Var}(X_1) = \sigma^2$ and $\mathbb{E}|X_1|^{2+\delta} < \infty$ for some $\delta > 0$, Davis and Mikosch (1999) shows that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j)/\sigma^2 - \log q \leq y\right) = \exp(-\exp(-y)). \quad (1.1)$$

The main purpose of this paper is to study the Cramér type moderate deviations for the maximum of the periodogram and its studentized version. That is, what is the largest possible a_n so that

$$\frac{\mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j)/\sigma^2 - \log q \geq y\right)}{1 - \exp(-\exp(-y))} \rightarrow 1 \quad (1.2)$$

uniformly in $y \in [-\log q, a_n]$, or for the studentized periodogram, what is the largest possible b_n so that

$$\frac{\mathbb{P}\left(\frac{\max_{1 \leq j \leq q} I_n(\omega_j)}{q^{-1} \sum_{j=1}^q I_n(\omega_j)} - \log q \geq y\right)}{1 - \exp(-\exp(-y))} \rightarrow 1 \quad (1.3)$$

uniformly in $y \in [-\log q, b_n]$. We shall show that a_n depends on the moment condition of $\{X_n\}$. For example, if $\mathbb{E}|X_1|^{2+\delta} < \infty$, $\delta > 0$, the largest possible value of a_n is $\frac{\delta}{2} \log n$, but a_n can be chosen $o(n^{1/3})$ if the moment generating function of X_1 is finite. However, the situation becomes totally different for the studentized periodogram. We shall prove that $b_n = o(n^{1/3})$ provided that $\mathbb{E}X_1^4 < \infty$.

The paper is organized as follows. Our main results, Theorems 2.1-2.3, are stated in Section 2, while proofs of the main results are postponed to Section 4. Our moderate deviation results are motivated by simultaneous tests in gene expression time series. Theoretical results for the simultaneous tests and simulation study are discussed in Section 3.

2 Main results

Throughout this paper, we assume $\{X_n\}$ are i.i.d. random variables. Our first result is the moderate deviation for the maximum of the periodogram for $y \leq c \log n$ for some $c > 0$. Such type of moderate deviation for the partial sums of $\{X_n\}$ has been studied in literature, e.g. by Michel (1976), Amosova (1982), Petrov (2002) and Wu and Zhao (2008).

Theorem 2.1 (i) Suppose that for some $c > 0$,

$$n^{c+1} \mathbb{P}\left(|X_1| \geq \sqrt{n \log n}\right) = o(1) \quad (2.1)$$

as $n \rightarrow \infty$. Then we have

$$\lim_{n \rightarrow \infty} \frac{P\left(\max_{1 \leq j \leq q} I_n(\omega_j)/\sigma^2 - \log q \geq y\right)}{1 - \exp(-\exp(-y))} = 1 \quad (2.2)$$

uniformly in $y \in [-\log q, c \log n]$, where $\sigma^2 = \text{Var}(X_1)$.

(ii) If for some $\sigma > 0$ (2.2) holds uniformly in $y \in [-\log q, c \log n]$ with some $c > 0$, then we have

$$n^{c+1}P\left(|X_1| \geq \sqrt{n \log n}\right) = O(1). \quad (2.3)$$

Theorem 2.1 (ii) shows that condition (2.1) is nearly optimal and hence the range depends on the moment assumption. On the other hand, when the moment generating function exists, the range can be extended to $o(n^{1/3})$.

Theorem 2.2 Assume $\text{Var}(X_1) = \sigma^2$ and $Ee^{t_0|X_1|} < \infty$ for some $t_0 > 0$. Then

$$\lim_{n \rightarrow \infty} \frac{P\left(\max_{1 \leq j \leq q} I_n(\omega_j)/\sigma^2 - \log q \geq y\right)}{1 - \exp(-\exp(-y))} = 1$$

uniformly in $y \in [-\log q, o(n^{1/3})]$.

We next consider the maximum of the studentized periodogram. Theorem 2.3 below shows that the moment conditions in Theorems 2.1 and 2.2 can be significantly reduced for the studentized version.

Theorem 2.3 If $EX_1^4 < \infty$, then

$$\lim_{n \rightarrow \infty} \frac{P\left(\frac{\max_{1 \leq j \leq q} I_n(\omega_j)}{q^{-1} \sum_{j=1}^q I_n(\omega_j)} - \log q \geq y\right)}{1 - \exp(-\exp(-y))} = 1 \quad (2.4)$$

uniformly in $y \in [-\log q, o(n^{1/3})]$.

Since the variance σ^2 of X_1 is typically unknown, what used in practice is actually the studentized periodogram. So the result in Theorem 2.3 is more appealing and useful than Theorems 2.1 and 2.2. Theorem 2.3 also shares similar properties with self-normalized partial sums of independent random variables, which usually requires much less moment assumptions; see, Shao (1997,1999) for self-normalized large deviation without any moment assumption and Cramér moderate deviation under finite third moment and de la Peña, Lai and Shao (2009) for recent developments in the area of self-normalized limit theory. In view of the moderate deviation for self-normalized partial sums (Shao (1999)), we conjecture that Theorem 2.3 remains true if $E|X_1|^3 < \infty$.

3 Application to simultaneous tests

3.1 Theoretical results

Periodic phenomena are widely studied in biology. Recently, there are quite a lot of interests in detecting periodic patterns in gene expression time series; see Wichert, Fokianos and Strimer (2004), Ahdesmäki et al. (2005), Chen (2005), Glynn, Chen and Mushegian (2006) and the references therein. Due to modern technology such as microarray experiments, the data are usually high-dimensional and we often need to make many statistical inference simultaneously. Let $Y_{t,g}$ denote the observed expression level of gene g at time t , $1 \leq g \leq G$ and $1 \leq t \leq n$, where G is the number of genes. The sample size n is usually much smaller than the number of genes. Consider the following model of periodic gene expression

$$Y_{t,g} = \mu_g + \beta_g \cos(\omega t + \phi) + \varepsilon_{t,g}, \quad (3.1)$$

where $\beta_g \geq 0$, $\omega \in (0, \pi)$, $\phi \in (-\pi, \pi]$, μ_g is the mean expression level. For each g , $\varepsilon_{1,g}, \dots, \varepsilon_{n,g}$ are i.i.d. noise sequence with mean zero. We wish to test the null hypothesis $H_{0,g} : \beta_g = 0$ against the alternative hypothesis $H_{1,g} : \beta_g \neq 0$. If $H_{0,g}$ is rejected, then we identify gene g with a periodic pattern in its expression. Periodogram is often used to detect periodically expressed gene. Let

$$I_n^{(g)}(\omega_j) = \frac{1}{n} \left| \sum_{k=1}^n Y_{k,g} e^{ik\omega_j} \right|^2, \quad (3.2)$$

where $\omega_j = 2\pi j/n$, $1 \leq j \leq [(n-1)/2]$. Define the g -statistic

$$f_g = \frac{\max_{1 \leq j \leq q} I_n^{(g)}(\omega_j)}{\sum_{j=1}^q I_n^{(g)}(\omega_j)},$$

and its null distribution $F_{n,g}(x) = P(f_g \leq x | H_{0,g})$, where $q = [(n-1)/2]$. Under the null hypothesis and the assumption that $\varepsilon_{1,g}, \dots, \varepsilon_{n,g}$ are i.i.d normal random variables, the exact distribution for f_g can be found in Fisher (1929):

$$P(f_g > x | H_{0,g}) = \sum_{j=1}^p (-1)^{j-1} C_q^j (1 - jx)^{q-1} =: f_n(x), \quad (3.3)$$

where $p = [1/x]$. Using (3.3), Wichert, Fokianos and Strimer (2004) proposed the following method to identify periodically expressed genes: 1. For each time series calculate Fisher's statistic f_g . 2. For each of the test statistic calculate the corresponding p value $P_g = f_n(f_g)$. 3. Use the method of Benjamini and Hochberg (1995)

to control the False Discovery Rate (FDR) at θ . Consider the set of ordered p values $P_{(1)}, \dots, P_{(G)}$ and let

$$i_\theta = \max\{i : P_{(i)} \leq i\theta/G\}. \quad (3.4)$$

Reject the null hypothesis for the time series indexed by $\mathcal{S} = \{i : P_i \leq P_{(i_\theta)}\}$.

In many applications such as those arising from bioinformatics, the noise can be remarkably non-Gaussian (Ahdesmäki et al. (2005)). Then the values $P_1, \dots, P_G$ are only the estimators of the true p -values. It is natural to ask:

How large G can be before the accuracy of simultaneous statistical inference becomes poor?

Similar problems have been studied in Fan, Hall and Yao (2007), where they consider $Y_{t,g} = \mu_g + \varepsilon_{t,g}$, the model in (3.1) without periodic parts, and focused on testing $H'_{0,g} : \mu_g = 0$. Let the true p value be $P_g^{true} = (1 - F_{n,g}(f_g))$. From (3.4), $P_{(i_\theta)}^{true}$ may be of the order $O(1/G)$. Hence $\max_{1 \leq g \leq G} |P_g^{true} - P_g| = o(1)$, implied by (1.1), is not enough. The required accuracy between the estimated p value and the true p value is

$$|P_g - P_g^{true}| I\{\mathcal{H}_g\} = o(P_g^{true}) \quad \text{uniformly in } 1 \leq g \leq G, \quad (3.5)$$

i.e.

$$\max_{1 \leq g \leq G} \left| \frac{P_g}{P_g^{true}} - 1 \right| I\{\mathcal{H}_g\} = o(1),$$

where $\mathcal{H}_g = \{P_g > \theta/(2G) \text{ or } P_g^{true} > \theta/(2G)\}$. (On $\mathcal{H}_g^c$, the gene g is always rejected.) Some similar requirements as (3.5) on simultaneous tests have been proposed by Fan, Hall and Yao (2007) and Kosorok and Ma (2007), p.1460.

Recall that $F_{n,g}(x) = \mathbf{P}(f_g \leq x | H_{0,g})$. By examining the proof of Theorem 2.3, we can get the following corollary.

Corollary 3.1 *Suppose that $\min_{1 \leq g \leq G} \text{Var}(\varepsilon_{1,g}) \geq \kappa$ for some $\kappa > 0$ which does not depend on G . Further assume that $\max_{1 \leq g \leq G} E\varepsilon_{1,g}^4 = O(1)$. Then the null distribution $F_{n,g}(x)$ satisfies*

$$\max_{1 \leq g \leq G} \left| \frac{1 - F_{n,g}((y + \log q)/q)}{1 - \exp(-\exp(-y))} - 1 \right| = o(1)$$

uniformly in $y \in [-\log q, o(n^{1/3})]$.

The following lemma shows that we can replace $1 - \exp(-\exp(-y))$ by $f_n((y + \log q)/q)$.

Lemma 3.1 *We have*

$$\lim_{n \rightarrow \infty} \left| \frac{f_n((y + \log q)/q)}{1 - \exp(-\exp(-y))} - 1 \right| = 0,$$

uniformly in $y \in [-\log q, o(n^{1/3})]$.

It follows from Corollary 3.1 and Lemma 3.1 that

Theorem 3.1 *Suppose the conditions in Corollary 3.1 are satisfied and $G = \exp(o(n^{1/3}))$. Then (3.5) holds.*

Theorem 3.1 shows that the level of the simultaneous tests is accurate provided that $G = \exp(o(n^{1/3}))$, which seems to be the correct order of asymptotics for microarray experiments with a moderate number of samples.

Using the bootstrap and a refined expansion of t -statistic in Theorem 1.2 of Wang (2005), Fan, Hall and Yao (2007) shows that $\exp(o(n^{1/3}))$ can be replaced by $\exp(o(n^{1/2}))$ for the tests $H_{0,g} : \mu_g = 0$. It would be interesting to investigate whether a similar expansion as Theorem 1.2 of Wang (2005) holds for the maximum of the studentized periodogram.

3.2 Simulation study

In this section, we carry out a simple simulation study to assess the finite sample performance. We generate 2000 genes with 100 periodic genes for different sample sizes n . Consider

$$\begin{aligned} Y_{t,g} &= \beta(\cos(\omega^{(g)}t) + \sin(\omega^{(g)}t)) + \varepsilon_{t,g}, & 1 \leq t \leq n, \quad 1 \leq g \leq 100, \\ Y_{t,g} &= \varepsilon_{t,g}, & 1 \leq t \leq n, \quad 101 \leq g \leq 2000. \end{aligned}$$

$\varepsilon_{t,g}$ will be taken as $N(0, 1)$, $(\sqrt{3/5}) \times t(5)$, $\text{EXP}(1)$, $2^{-1} \times \chi^2(2)$, where $t(5)$ has the t distribution with freedom 5, $\text{EXP}(1)$ is the exponential random variable with parameter $\mu = 1$, $\chi^2(2)$ is chi square random variable with freedom 2. (The constants on the left hand side of random variables are chosen so that $\text{Var}(\varepsilon_{t,g}) = 1$). The FDR level θ is chosen as 0.15 and 0.05. The simulation results are based on 100 replicates.

We only give the simulation study when $\omega^{(g)}$ is of the form of ω_i for some $1 \leq i \leq q$. To do this, we let $\omega^{(g)} = 2\pi/10$, $n = 20, 50$ and $\beta = 1$. The results are summarized in Table 1, where Tot.=total count identified using FDR; Pos.= the number of true positives identified using FDR; Z=the number of true periodic genes among the smallest 100 p -values genes. We note that when the tails of $\varepsilon_{t,g}$ are heavier

than that of Gaussian random variable, the empirical FDR (EFDR) are lower than the target FDR, while most of periodic genes can still be found. There are no significant differences between Gaussian noise and other noises when n is large moderately ($n = 50$). Powers increase as n increases. Overall Fisher's statistic is relatively robust to the noise, as indicated by our theorem. We refer to Wichert, Fokianos and Strimer (2004) for some real data analysis.

Table 1:

θ		Tot.(Pos.)	EFDR	Tot.(Pos.)	EFDR
		$n = 20$		$n = 50$	
0.15	Normal	39.9(33.7)	0.155	116.2(99.6)	0.143
0.05		12.6(11.9)	0.059	102.6(99.2)	0.033
Z		62		97.1	
0.15	EXP(1)	55.3(49.5)	0.105	104.3(97.3)	0.067
0.05		33.1(31.9)	0.036	97.8(95.8)	0.020
Z		69.5		96.9	
0.15	$\chi^2(2)$	46.0(43.2)	0.061	105.0(97.6)	0.071
0.05		29.0(28.2)	0.028	98.1(95.6)	0.026
Z		69.7		96.5	
0.15	$t(5)$	43.4(39.7)	0.085	110.2(98.2)	0.109
0.05		20.2(19.6)	0.030	99.9(96.6)	0.033
Z		67.6		96.6	

4 Proofs

Throughout, we let C denote positive constant, and its value may be different in different contexts. For two real sequences $\{a_n\}$ and $\{b_n\}$, write $a_n = O(b_n)$ if there exists a constant C such that $|a_n| \leq C|b_n|$ holds for large n , $a_n = o(b_n)$ if $\lim_{n \rightarrow \infty} a_n/b_n = 0$. We denote by $|\cdot|$ the d -dimensional Euclidean norm in $\mathbf{R}^d$, $d \geq 1$.

Proof of Theorem 2.1. (i) (**Sufficiency**) By $\sum_{k=1}^n e^{ik\omega_j} = 0$ for $1 \leq j \leq q$, we can assume that $\mathbf{E}X_1 = 0$. Also, for convenience, we assume $\sigma^2 = 1$. For $y \in [-\log q, c \log n]$, set $x = \sqrt{y + \log q}$. We start with truncation of X_k at two levels. Let $\varepsilon_n = (\log n)^{-1}$ and $\varepsilon > 0$ be a small number which will be specified later. Define

$$\begin{aligned}
\tilde{X}_k &= X'_k - \mathbf{E}X'_k, & X'_k &= X_k I\{|X_k| \leq \varepsilon \sqrt{nx}\}, \\
\hat{X}_k &= X''_k - \mathbf{E}X''_k, & X''_k &= X_k I\{|X_k| \leq \varepsilon_n \sqrt{n}/x\}, \\
I'_n(\omega_j) &= \frac{1}{n} \left| \sum_{k=1}^n X'_k e^{ik\omega_j} \right|^2 = \frac{1}{n} \left| \sum_{k=1}^n \tilde{X}_k e^{ik\omega_j} \right|^2,
\end{aligned}$$

$$I_n''(\omega_j) = \frac{1}{n} \left| \sum_{k=1}^n X_k'' e^{ik\omega_j} \right|^2 = \frac{1}{n} \left| \sum_{k=1}^n \widehat{X}_k e^{ik\omega_j} \right|^2. \quad (4.1)$$

Then we have

$$\begin{aligned} & \left| \mathbb{P} \left(\max_{1 \leq j \leq q} I_n(\omega_j) \geq x^2 \right) - \mathbb{P} \left(\max_{1 \leq j \leq q} I_n'(\omega_j) \geq x^2 \right) \right| \\ & \leq n \mathbb{P} \left(|X_1| \geq \varepsilon \sqrt{n} x \right). \end{aligned} \quad (4.2)$$

The independence between X_l and $\{X_k, k \neq l\}$, $1 \leq l \leq n$, implies that

$$\begin{aligned} & \mathbb{P} \left(\max_{1 \leq j \leq q} I_n'(\omega_j) \geq x^2 \right) - \mathbb{P} \left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2 \right) \\ & \leq \mathbb{P} \left(\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k' e^{ik\omega_j} \right| \geq \sqrt{n} x, \cup_{l=1}^n \{|X_l| > \varepsilon_n \sqrt{n}/x\} \right) \\ & \leq \sum_{l=1}^n \mathbb{P} \left(\max_{1 \leq j \leq q} \left| \sum_{k=1, k \neq l}^n X_k' e^{ik\omega_j} \right| \geq (1 - \varepsilon) \sqrt{n} x \right) \mathbb{P} \left(|X_l| > \varepsilon_n \sqrt{n}/x \right) \\ & \leq n \mathbb{P} \left(\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k' e^{ik\omega_j} \right| \geq (1 - 2\varepsilon) \sqrt{n} x \right) \mathbb{P} \left(|X_1| > \varepsilon_n \sqrt{n}/x \right) \\ & \leq n \mathbb{P} \left(\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k'' e^{ik\omega_j} \right| \geq (1 - 2\varepsilon) \sqrt{n} x \right) \mathbb{P} \left(|X_1| > \varepsilon_n \sqrt{n}/x \right) \\ & \quad + \left(n \mathbb{P} \left(|X_1| > \varepsilon_n \sqrt{n}/x \right) \right)^2 \\ & =: A + \left(n \mathbb{P} \left(|X_1| > \varepsilon_n \sqrt{n}/x \right) \right)^2. \end{aligned} \quad (4.3)$$

To estimate A , we need Lemma 4.2 of Lin and Liu (2009a). The proof is given in Lin and Liu (2009b), pages 23-25 and the constants n_0 , C , $c_{1,1}, \dots, c_{1,3}$ below are specified in Lin and Liu (2009b).

Lemma 4.1 [Lin and Liu (2009a,b)] Let $\xi_{n,1}, \dots, \xi_{n,k_n}$ be independent random vectors with mean zero and values in $\mathbf{R}^{2d}$, and $S_n = \sum_{i=1}^{k_n} X_{n,i}$. Assume that $|\xi_{n,k}| \leq c_n B_n^{1/2}$, $1 \leq k \leq k_n$, for some $c_n \rightarrow 0$, $B_n \rightarrow \infty$ and

$$\left\| B_n^{-1} \text{Cov}(\xi_{n,1} + \dots + \xi_{n,k_n}) - I_{2d} \right\| \leq C_0 c_n^2,$$

where I_{2d} is a $2d \times 2d$ identity matrix and C_0 is a positive constant. Suppose that $\beta_n := B_n^{-3/2} \sum_{k=1}^{k_n} E|\xi_{n,k}|^3 \rightarrow 0$. Then for all $n \geq n_0$ (n_0 is given below)

$$\begin{aligned} & |P(\|S_n\|_d \geq x) - P(\|N\|_d \geq x/B_n^{1/2})| \\ & \leq o(1)P(\|N\|_d \geq x/B_n^{1/2}) + C \left(\exp \left(- \frac{\delta_n^2 \min(c_n^{-2}, \beta_n^{-2/3})}{16d} \right) + \exp \left(\frac{C c_n^2}{\beta_n^2 \log \beta_n} \right) \right), \end{aligned}$$

uniformly for $x \in [B_n^{1/2}, \delta_n \min(c_n^{-1}, \beta_n^{-1/3}) B_n^{1/2}]$, with any $\delta_n \rightarrow 0$ and $\delta_n \min(c_n^{-1}, \beta_n^{-1/3}) \rightarrow \infty$. N is a centered normal random vector with covariance matrix I_{2d} . $\|\cdot\|_d$ is defined

by $\|z\|_d = \min\{(x_i^2 + y_i^2)^{1/2} : 1 \leq i \leq d\}$, $z = (x_1, y_1, \dots, x_d, y_d)$. $\|\cdot\|$ is the spectral norm for the matrix. C is a positive constant which depends only on d and C_0 . $o(1)$ is bounded by $c_{1,1}(\delta_n^3 + \beta_n + c_n)$, $c_{1,1}$ is a positive constant depending only on d .

$$n_0 = \min \left\{ n : \forall k \geq n, \ c_k^2 \leq \frac{\min(C_0^{-1}, 8^{-1})}{2}, \ \delta_k \leq c_{1,2} \min(C_0^{-2}, 1), \ \beta_k \leq c_{1,3} \right\},$$

where $c_{1,2}$ and $c_{1,3}$ are positive constants depending only on d .

Let $d \geq 1$ be a fixed integer and

$$\begin{aligned} \mathbf{Y}_k &:= \mathbf{Y}_k(\omega_{i_1}, \dots, \omega_{i_d}) \\ &= \widehat{X}_k \left(\cos(k\omega_{i_1}), \sin(k\omega_{i_1}), \dots, \cos(k\omega_{i_d}), \sin(k\omega_{i_d}) \right), \\ 1 \leq k \leq n, \quad 1 \leq i_1 < \dots < i_d \leq q. \end{aligned}$$

Let $\|z\|_d = \min\{(x_k^2 + y_k^2)^{1/2} : 1 \leq k \leq d\}$ with $z = (x_1, y_1, \dots, x_d, y_d)$. By the facts that for $1 \leq j, l \leq q$,

$$\sum_{k=1}^n \cos^2(\omega_j k) = n/2, \quad \sum_{k=1}^n \sin^2(\omega_j k) = n/2, \quad \sum_{k=1}^n \cos(\omega_j k) \sin(\omega_l k) = 0, \quad (4.4)$$

we can see that

$$\left\| n^{-1} \text{Cov} \left(\sum_{k=1}^n \mathbf{Y}_k \right) - \frac{1}{2} \mathbf{I}_{2d} \right\| \leq \mathbf{E} X_1^2 I\{|X_1| \geq \varepsilon_n \sqrt{n}/x\}, \quad (4.5)$$

where $\mathbf{I}_{2d}$ is $2d \times 2d$ identity matrix and $\|\cdot\|$ is the spectral norm. It is easy to see that (2.1) implies $\mathbf{E}|X_1|^p < \infty$ for any $2 < p < 2c + 2$. Thus we have

$$|\mathbf{Y}_k| \leq 2d\varepsilon_n \sqrt{n}/x \text{ and } n^{-3/2} \sum_{k=1}^n \mathbf{E}|\mathbf{Y}_k|^3 \leq C \max(n^{1-p/2} x^{-3+p}, n^{-1/2}). \quad (4.6)$$

Letting $c_n = 2\varepsilon_n(\log q)^{-1/2}$, $B_n = n$ in Lemma 4.1 and $\delta_n \log n \rightarrow \infty$ with δ_n being defined in Lemma 4.1, we have for any $0 \leq \eta < 1$,

$$\frac{\mathbf{P} \left(\left\| \sum_{k=1}^n \mathbf{Y}_k \right\|_d \geq (1 - \eta) \sqrt{nx} \right)}{q^{-d(1-\eta)^2} \exp(-d(1-\eta)^2 y)} \rightarrow 1, \quad (4.7)$$

uniformly in $x \in [\sqrt{\log q}, \sqrt{(c+1) \log n}]$. Observing that

$$I_n''(\omega_j) = \frac{1}{n} \left(\left\| \sum_{k=1}^n \mathbf{Y}_k(\omega_j) \right\|_1 \right)^2,$$

by (4.7),

$$\mathbf{P} \left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2 \right) \leq \sum_{j=1}^q \mathbf{P} \left(\left\| \sum_{k=1}^n \mathbf{Y}_k(\omega_j) \right\|_1 \geq \sqrt{nx} \right) \leq (1 + o(1)) \exp(-y)$$

uniformly in $y \in [0, c \log n]$. So, combining (4.2)-(4.7), we prove that

$$\begin{aligned} \mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) \geq x^2\right) &\leq (1 + o(1)) \exp(-y) \\ &\quad + Cn^{1+4\varepsilon} \exp(-(1-2\varepsilon)^2 y) \mathbb{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right) \\ &\quad + \left(n \mathbb{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right)\right)^2 + n \mathbb{P}\left(|X_1| \geq \varepsilon \sqrt{n} x\right), \end{aligned} \quad (4.8)$$

uniformly in $x \in [\sqrt{\log q}, \sqrt{(c+1) \log n}]$. To establish the lower bound, we observe that

$$\begin{aligned} \mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) \geq x^2\right) &\geq \mathbb{P}\left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2, \cap_{k=1}^n \{|X_k| \leq \varepsilon_n \sqrt{n}/x\}\right) \\ &= \mathbb{P}\left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2\right) \\ &\quad - \mathbb{P}\left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2, \cup_{k=1}^n \{|X_k| \geq \varepsilon_n \sqrt{n}/x\}\right). \end{aligned} \quad (4.9)$$

Similarly to (4.3) and by (4.7), we have

$$\begin{aligned} &\mathbb{P}\left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2, \cup_{k=1}^n \{|X_k| \geq \varepsilon_n \sqrt{n}/x\}\right) \\ &\leq n \mathbb{P}\left(\max_{1 \leq j \leq q} \left|\sum_{k=1}^n X_k'' e^{ik\omega_j}\right| \geq (1-2\varepsilon)\sqrt{n}x\right) \mathbb{P}(|X_1| \geq \varepsilon_n \sqrt{n}/x) \\ &\leq Cn^{1+4\varepsilon} \exp(-(1-2\varepsilon)^2 y) \mathbb{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right) \end{aligned} \quad (4.10)$$

uniformly in $x \in [\sqrt{\log q}, \sqrt{(c+1) \log n}]$. For the first term in (4.9), we have

$$\mathbb{P}\left(\max_{1 \leq j \leq q} I_n''(\omega_j) \geq x^2\right) \geq \sum_{j=1}^q \mathbb{P}(A_j) - \sum_{1 \leq i < j \leq q} \mathbb{P}(A_i A_j), \quad (4.11)$$

where $A_j = \{I_n''(\omega_j) \geq x^2\}$. Applying $d = 1, 2$ in (4.7) respectively, we obtain

$$\mathbb{P}(A_i) = (1 + o(1))q^{-1} \exp(-y), \quad \mathbb{P}(A_i A_j) \leq Cn^{-2} \exp(-2y),$$

uniformly in $1 \leq i, j \leq q$ and $y \in [0, c \log n]$. These two inequalities together with (4.9)-(4.11) yield that

$$\begin{aligned} \mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) \geq x^2\right) &\geq (1 + o(1)) \exp(-y) - C \exp(-2y) \\ &\quad - Cn^{1+4\varepsilon} \exp(-(1-2\varepsilon)^2 y) \mathbb{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right), \end{aligned} \quad (4.12)$$

uniformly in $y \in [0, c \log n]$. It is easy to see that (2.1) implies that

$$\mathbb{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right) \leq C \frac{(\log n)^{4c+4}}{n^{c+1}}$$

for $\varepsilon_n = (\log n)^{-1}$ and $x \in [\sqrt{\log q}, \sqrt{(c+1)\log n}]$. Hence, by (4.8), (4.12) and for ε sufficiently small, we have for any $M > 0$,

$$\limsup_{n \rightarrow \infty} \sup_{M \leq y \leq c \log n} \left| \frac{\mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) - \log q \geq y\right)}{1 - \exp(-\exp(-y))} - 1 \right| \leq Ce^{-M}. \quad (4.13)$$

By (1.1), we have for any fixed $y \in \mathbf{R}$,

$$\mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) - \log q \geq y\right) \rightarrow 1 - e^{-e^{-y}}.$$

Thus, it follows that

$$\limsup_{n \rightarrow \infty} \sup_{-\log q \leq y \leq M} \left| \frac{\mathbb{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) - \log q \geq y\right)}{1 - \exp(-\exp(-y))} - 1 \right| = 0. \quad (4.14)$$

This proves (i) by (4.13) and (4.14).

(ii) **(Necessity)** By (2.2), we have

$$\mathbb{P}\left(\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k e^{ik\omega_j} \right| \geq \sigma \sqrt{(1+c)n \log q}\right) \leq Cn^{-c}.$$

This implies that

$$\mathbb{P}\left(\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k^s e^{ik\omega_j} \right| \geq 2^{-1} \sigma \sqrt{(1+c)n \log q}\right) \leq Cn^{-c}, \quad (4.15)$$

where $X_n^s = X_n - X_n^c$ and $\{X_n^c\}$ is an independent copy of $\{X_n\}$. For $z = (z_{1,1}, z_{1,2}, \dots, z_{q,1}, z_{q,2}) \in \mathbf{R}^{2q}$, let

$$\|z\|_{\max} = \max_{1 \leq j \leq q} \sqrt{z_{j,1}^2 + z_{j,2}^2}.$$

For $1 \leq k \leq n$, let

$$\mathbf{D}_k = (X_k^s \cos(k\omega_1), X_k^s \sin(k\omega_1), \dots, X_k^s \cos(k\omega_q), X_k^s \sin(k\omega_q)).$$

Then it is easy to see that $\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k^s e^{ik\omega_j} \right| = \left\| \sum_{k=1}^n \mathbf{D}_k \right\|_{\max}$. By Lévy's inequality in a Banach space (cf. Ledoux and Talagrand (1991), p. 47) and (4.15), we have

$$\mathbb{P}\left(\max_{1 \leq k \leq n} \|\mathbf{D}_k\|_{\max} \geq 2^{-1} \sigma \sqrt{(1+c)n \log q}\right) \leq Cn^{-c}. \quad (4.16)$$

Observing that $\|\mathbf{D}_k\|_{\max} = |X_k^s|$, we have by (4.16),

$$1 - \left(1 - \mathbb{P}\left(|X_1^s| \geq 2^{-1} \sigma \sqrt{(1+c)n \log q}\right)\right)^n \leq Cn^{-c},$$

which implies that

$$1 - \exp\left(-n\mathbf{P}\left(|X_1^s| \geq 2^{-1}\sigma\sqrt{(1+c)n\log q}\right)\right) \leq Cn^{-c}. \quad (4.17)$$

Now (2.3) follows from (4.17) and some elementary calculations. $\square$

Proof of Theorem 2.2. For any $\varepsilon'_n \rightarrow 0$, let $x = \sqrt{y + \log q} \in [4, \varepsilon'_n n^{1/6}]$ and $\varepsilon_n = \varepsilon'_n{}^{1/4}$. Define

$$X'_k = X_k I\{|X_k| \leq \varepsilon_n \sqrt{n}/x\}, \quad I'_n(\omega_j) = \frac{1}{n} \left| \sum_{k=1}^n X'_k e^{ik\omega_j} \right|^2.$$

Then, by Lemma 4.2 below (taking $\mathbf{N}_l = \emptyset$ and $d = 1$), we have

$$\mathbf{P}\left(\max_{1 \leq j \leq q} I'_n(\omega_j) \geq x^2\right) \leq \sum_{j=1}^q \mathbf{P}\left(\left| \sum_{k=1}^n X'_k e^{ik\omega_j} \right| \geq \sqrt{nx}\right) \leq (1 + o(1))e^{-y}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. Since $\mathbf{E}e^{t_0|X_1|} < \infty$, we have $n\mathbf{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right) = o(1)e^{-x^2}$. Therefore,

$$\begin{aligned} \mathbf{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) \geq x^2\right) &\leq \mathbf{P}\left(\max_{1 \leq j \leq q} I'_n(\omega_j) \geq x^2\right) + n\mathbf{P}\left(|X_1| > \varepsilon_n \sqrt{n}/x\right) \\ &\leq (1 + o(1))e^{-y} \end{aligned}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. Similarly, we have

$$\mathbf{P}\left(\max_{1 \leq j \leq q} I_n(\omega_j) \geq x^2\right) \geq (1 + o(1))e^{-y} - Ce^{-2y}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. The remaining proof follows similar arguments as in the proof of Theorem 2.1. $\square$

Proof of Theorem 2.3. Recall that $\sum_{k=1}^n e^{ik\omega_j} = 0$ for $1 \leq j \leq q$. Without loss of generality, we can assume that $\mathbf{E}X_1 = 0$ and $\mathbf{E}X_1^2 = 1$. By the fact $e^{ik\omega_{j_1}} = e^{-ik\omega_{j_2}}$ for $j_1 + j_2 = n$, we have

$$\left| \sum_{k=1}^n X_k e^{ik\omega_{j_1}} \right|^2 = \left| \sum_{k=1}^n X_k e^{ik\omega_{j_2}} \right|^2$$

for $j_1 + j_2 = n$. Hence, when n is odd,

$$\begin{aligned} q^{-1} \sum_{j=1}^q I_n(\omega_j) &= (2q)^{-1} \sum_{j=1}^q \left(I_n(\omega_j) + I_n(\omega_{n-j}) \right) \\ &= (2q)^{-1} \sum_{j=1}^n I_n(\omega_j) - \frac{n}{n-1} (\overline{X})^2, \end{aligned} \quad (4.18)$$

where $\overline{X} = n^{-1} \sum_{k=1}^n X_k$. Moreover,

$$\sum_{j=1}^n I_n(\omega_j) = \sum_{k=1}^n X_k^2 + 2n^{-1} \sum_{k=2}^n X_k \sum_{i=1}^{k-1} X_i \sum_{j=1}^n w_{k,i,j},$$

where $w_{k,i,j} = \cos(\omega_j k) \cos(\omega_j i) + \sin(\omega_j k) \sin(\omega_j i)$. Note that $\omega_j k = \omega_k j$ and $\omega_j i = \omega_i j$. Since $|\sum_{l=1}^n e^{il\lambda}| = |\sin(\lambda n/2)|/|\sin(\lambda/2)|$ when λ/π is not an integer, we get $\sum_{j=1}^n w_{k,i,j} = \sum_{j=1}^n \cos((\omega_k - \omega_i)j) = 0$. So, (4.18) implies that, when n is odd,

$$q^{-1} \sum_{j=1}^q I_n(\omega_j) = (n-1)^{-1} \left(\sum_{k=1}^n X_k^2 - n(\overline{X})^2 \right).$$

Similarly, when n is even, we have

$$q^{-1} \sum_{j=1}^q I_n(\omega_j) = (n-2)^{-1} \left(\sum_{k=1}^n X_k^2 - n(\overline{X})^2 - n(\overline{X}')^2 \right), \quad (4.19)$$

where $\overline{X}' = n^{-1} \sum_{k=1}^n (-1)^k X_k$. By the self-normalized moderate deviation Theorem 3.1 in Shao (1997), we have

$$\mathbb{P}\left(\left|\sum_{k=1}^n (-1)^k X_k\right| \geq n^{1/3} V_n\right) \leq C e^{-n^{2/3}/4}, \quad \mathbb{P}\left(\left|\sum_{k=1}^n X_k\right| \geq n^{1/3} V_n\right) \leq C e^{-n^{2/3}/4},$$

where $V_n^2 = \sum_{k=1}^n X_k^2$. In view of (4.18) and (4.19), it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{\mathbb{P}\left(\frac{\max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k e^{ik\omega_j} \right|^2}{\sum_{k=1}^n X_k^2} - \log q \geq y\right)}{1 - e^{-e^{-y}}} = 1 \quad (4.20)$$

uniformly in $y \in [-\log q, o(n^{1/3})]$.

Let $\lambda = \lambda_n$ be a positive number which will be specified later. Let $\mathbf{H}$ be a subset of $\{1, \dots, n\}$. Put

$$\begin{aligned} X'_k &= X_k I\{|X_k| \leq \lambda\}, \quad 1 \leq k \leq n, \\ M_n &= \max_{1 \leq j \leq q} \left| \sum_{k=1}^n X_k e^{ik\omega_j} \right|, \quad \widetilde{M}_n = \max_{1 \leq j \leq q} \left| \sum_{k=1}^n X'_k e^{ik\omega_j} \right|, \\ M_n^{(\mathbf{H})} &= \max_{1 \leq j \leq q} \left| \sum_{k=1, k \notin \mathbf{H}}^n X_k e^{ik\omega_j} \right|, \quad \widetilde{M}_n^{(\mathbf{H})} = \max_{1 \leq j \leq q} \left| \sum_{k=1, k \notin \mathbf{H}}^n X'_k e^{ik\omega_j} \right|, \\ \widetilde{V}_n &= \left(\sum_{k=1}^n X_k'^2 \right)^{1/2}, \quad V_n^{(\mathbf{H})} = \left(\sum_{k=1, k \notin \mathbf{H}}^n X_k^2 \right)^{1/2}, \quad \widetilde{V}_n^{(\mathbf{H})} = \left(\sum_{k=1, k \notin \mathbf{H}}^n X_k'^2 \right)^{1/2}. \end{aligned}$$

Noting that for any real numbers s and t and nonnegative number c and $x \geq 1$,

$$\{s + t \geq x\sqrt{c + t^2}\} \subset \{s \geq \sqrt{x^2 - 1}\sqrt{c}\}, \quad (4.21)$$

(see p.2181 in Jing, Shao and Wang (2003)), we have

$$\begin{aligned}
\mathbb{P}(M_n \geq xV_n) &\leq \mathbb{P}(\widetilde{M}_n \geq x\widetilde{V}_n) + \sum_{j=1}^n \mathbb{P}(M_n \geq xV_n, X_j \neq X'_j) \\
&\leq \mathbb{P}(\widetilde{M}_n \geq x\widetilde{V}_n) + \sum_{j=1}^n \mathbb{P}(M_n^{(j)} + |X_j| \geq xV_n, X_j \neq X'_j) \\
&\leq \mathbb{P}(\widetilde{M}_n \geq x\widetilde{V}_n) + \sum_{j=1}^n \mathbb{P}(M_n^{(j)} \geq \sqrt{x^2 - 1}V_n^{(j)}, X_j \neq X'_j) \\
&= \mathbb{P}(\widetilde{M}_n \geq x\widetilde{V}_n) + \sum_{j=1}^n \mathbb{P}(|X_j| \geq \lambda) \mathbb{P}(M_n^{(j)} \geq \sqrt{x^2 - 1}V_n^{(j)}) \quad (4.22)
\end{aligned}$$

Repeating the above arguments m times with $m = \lfloor x^2/2 \rfloor$, we have for $x > 4$,

$$\begin{aligned}
&\sum_{j_1=1}^n \mathbb{P}(|X_{j_1}| \geq \lambda) \mathbb{P}(M_n^{(j_1)} \geq \sqrt{x^2 - 1}V_n^{(j_1)}) \\
&\leq \sum_{j_1=1}^n \mathbb{P}(|X_{j_1}| \geq \lambda) \mathbb{P}(\widetilde{M}_n^{(j_1)} \geq \sqrt{x^2 - 1}\widetilde{V}_n^{(j_1)}) \\
&\quad + \sum_{j_1=1}^n \sum_{j_2=1}^n \mathbb{P}(|X_{j_1}| \geq \lambda) \mathbb{P}(|X_{j_2}| \geq \lambda) \mathbb{P}(M_n^{(j_1, j_2)} \geq \sqrt{x^2 - 2}V_n^{(j_1, j_2)}) \\
&\leq \sum_{l=1}^m \widetilde{Z}_l + Z_{m+1}, \quad (4.23)
\end{aligned}$$

where

$$\begin{aligned}
\widetilde{Z}_l &= \sum_{j_1=1}^n \cdots \sum_{j_l=1}^n \left[\prod_{k=1}^l \mathbb{P}(|X_{j_k}| \geq \lambda) \right] \times \mathbb{P}(\widetilde{M}_n^{(j_1, \dots, j_l)} \geq \sqrt{x^2 - l}\widetilde{V}_n^{(j_1, \dots, j_l)}), \\
Z_{m+1} &= \sum_{j_1=1}^n \cdots \sum_{j_{m+1}=1}^n \left\{ \left[\prod_{k=1}^{m+1} \mathbb{P}(|X_{j_k}| \geq \lambda) \right] \right. \\
&\quad \left. \times \mathbb{P}(M_n^{(j_1, \dots, j_{m+1})} \geq \sqrt{x^2 - m - 1}V_n^{(j_1, \dots, j_{m+1})}) \right\}.
\end{aligned}$$

For $\varepsilon'_n \rightarrow 0$ and $4 \leq x \leq \varepsilon'_n n^{1/6}$, let $\lambda = \varepsilon_n \sqrt{n}/x$, where $\varepsilon_n = \varepsilon_n'^{1/4}$. Then

$$Z_{m+1} \leq \left(n \mathbb{P}(|X_1| \geq \lambda) \right)^{m+1} \leq e^{-m \log q_n} = o(1) e^{-x^2}, \quad (4.24)$$

where

$$q_n = \left(\varepsilon_n'^3 \varepsilon_n^{-3} \mathbb{E}|X_1|^3 I\{|X_1| \geq \lambda\} \right)^{-1} \rightarrow \infty$$

as $n \rightarrow \infty$. Let $0 \leq l \leq m = \lfloor x^2/2 \rfloor$ and $\mathbf{N}_l = \{j_1, \dots, j_l\} \subset \{1, \dots, n\}$. Define

$$\mathbf{Y}'_k = X'_k \left(\cos(k\omega_{i_1}), \sin(k\omega_{i_1}), \dots, \cos(k\omega_{i_d}), \sin(k\omega_{i_d}) \right),$$

$$1 \leq k \leq n, \quad d \geq 1, \quad 1 \leq i_1 < \cdots < i_d \leq q.$$

$$S_n^{\mathbf{N}_l} = \sum_{k=1, k \notin \mathbf{N}_l}^n \mathbf{Y}'_k, \quad \bar{S}_n^{\mathbf{N}_l} = \sum_{k=1, k \notin \mathbf{N}_l}^n (\mathbf{Y}'_k - \mathbf{E}\mathbf{Y}'_k).$$

To estimate $\tilde{Z}_l$, we need the following lemma.

Lemma 4.2 *Suppose that $\mathbf{E}|X_1|^3 < \infty$ and $0 \leq x \leq \varepsilon'_n n^{1/6}$, where $\varepsilon'_n \rightarrow 0$ is any sequence of constants. Let $0 < \varepsilon_n \rightarrow 0$ and $\varepsilon_n = \varepsilon_n'^{1/4}$. (i) If $\lambda = \varepsilon_n \sqrt{n}/x$, then we have*

$$\lim_{n \rightarrow \infty} \frac{P\left(\left\|S_n^{\mathbf{N}_l}\right\|_d \geq x\sqrt{n}\right)}{e^{-dx^2}} = 1 \quad (4.25)$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$, $1 \leq i_1 < \cdots < i_d \leq q$ and $0 \leq l \leq m$. (ii) If $\lambda = (\varepsilon_n \sqrt{n}/x)^{3/4}$ and $\mathbf{E}X_1^4 < \infty$, then (4.25) holds.

Proof. Recall that $\sum_{k=1}^n e^{ik\omega_j} = 0$ for $1 \leq j \leq q$. We have for n large,

$$\left| \sum_{k=1, k \notin \mathbf{N}_l}^n \mathbf{E}\mathbf{Y}'_k \right| = \left| \sum_{k \in \mathbf{N}_l}^n \mathbf{E}\mathbf{Y}'_k \right| \leq dx^2/2 \leq \varepsilon_n \sqrt{n}/x.$$

It follows that, for $x \in [4, \varepsilon'_n n^{1/6}]$,

$$\begin{aligned} P\left(\left\|\bar{S}_n^{\mathbf{N}_l}\right\|_d \geq x\sqrt{n} + \varepsilon_n \sqrt{n}/x\right) &\leq P\left(\left\|S_n^{\mathbf{N}_l}\right\|_d \geq x\sqrt{n}\right) \\ &\leq P\left(\left\|\bar{S}_n^{\mathbf{N}_l}\right\|_d \geq x\sqrt{n} - \varepsilon_n \sqrt{n}/x\right). \end{aligned}$$

Since $\mathbf{E}|X_1|^3 < \infty$, we have

$$|\mathbf{Y}'_k| \leq 2d\varepsilon_n \sqrt{n}/x \text{ and } n^{-3/2} \sum_{k=1}^n \mathbf{E}|\mathbf{Y}_k|^3 \leq Cn^{-1/2}.$$

Also, simple calculations show that

$$\left\| \frac{1}{n} \text{Cov}\left(\bar{S}_n^{\mathbf{N}}\right) - \mathbf{I}_{2d} \right\| \leq \mathbf{E}X_1^2 I\{|X_1| \geq \lambda\} + C_d n^{-1} x^2 \leq Cx n^{-1/2} \varepsilon_n^{-1} \leq C\varepsilon_n^2/x^2$$

for $4 \leq x \leq \varepsilon'_n n^{1/6}$. By taking $c_n = 2d\varepsilon_n/x$, $B_n = \sqrt{n}$ and $\delta_n/\varepsilon_n \rightarrow \infty$ in Lemma 4.1, we have

$$\frac{P\left(\left\|\bar{S}_n^{\mathbf{N}_l}\right\|_d \geq x\sqrt{n} \pm \varepsilon_n \sqrt{n}/x\right)}{e^{-dx^2}} \rightarrow 1,$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. This proves Lemma 4.2.

From Lemma 4.2, we have for $0 \leq l \leq m = \lfloor x^2/2 \rfloor$,

$$P\left(\widetilde{M}_n^{(j_1, \dots, j_l)} \geq \sqrt{x^2 - l} \widetilde{V}_n^{(j_1, \dots, j_l)}\right)$$

$$\begin{aligned}
&\leq \mathbb{P}\left(\widetilde{M}_n^{(j_1, \dots, j_l)} \geq \sqrt{x^2 - l} \sqrt{n(1 - \varepsilon_n x^{-2})}\right) \\
&\quad + \mathbb{P}\left(\widetilde{V}_n^{(j_1, \dots, j_l)} \leq \sqrt{n(1 - \varepsilon_n x^{-2})}\right) \\
&= (1 + o(1)) q e^{-x^2 + l} + \mathbb{P}\left(\widetilde{V}_n^{(j_1, \dots, j_l)} \leq \sqrt{n(1 - \varepsilon_n x^{-2})}\right), \tag{4.26}
\end{aligned}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. By a similar argument as in Hu, Shao and Wang (2009), p. 1193, if $\mathbb{E}X_1^4 < \infty$, we have

$$\mathbb{P}\left(\widetilde{V}_n^{(j_1, \dots, j_l)} \leq \sqrt{n(1 - \varepsilon_n x^{-2})}\right) \leq \mathbb{P}\left(n - m_l - \widetilde{V}_n^{(j_1, \dots, j_l)2} \geq \varepsilon_n n x^{-2}/2\right) \leq o(1) e^{-x^2},$$

where m_l is the cardinality of $\{j_1, \dots, j_l\}$, and hence

$$\widetilde{Z}_l \leq Cn \left(n \mathbb{P}\left(|X_1| \geq \lambda\right) \right)^l e^{-(x^2 - l)} \leq Cn e^{-x^2 + l - Cl \log q_n}. \tag{4.27}$$

This together with $q_n \rightarrow \infty$ shows that

$$\sum_{l=1}^m \widetilde{Z}_l = o(1) n e^{-x^2}, \tag{4.28}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. Combining (4.22)-(4.24), (4.26) and (4.28) yields

$$\mathbb{P}\left(M_n \geq x V_n\right) \leq (1 + o(1)) q e^{-x^2} \tag{4.29}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$.

We next estimate the lower bound for $\mathbb{P}\left(M_n \geq x V_n\right)$. For $\varepsilon'_n \rightarrow 0$ and $4 \leq x \leq \varepsilon'_n n^{1/6}$, let $\varepsilon_n = \max((x/\sqrt{n})^{1/8}, \varepsilon_n'^{1/4})$ and $\lambda = (\varepsilon_n \sqrt{n}/x)^{3/4}$. Then

$$\begin{aligned}
\mathbb{P}\left(M_n \geq x V_n\right) &\geq \mathbb{P}\left(\widetilde{M}_n \geq x \widetilde{V}_n\right) \\
&\quad - \sum_{j=1}^n \mathbb{P}\left(\widetilde{M}_n^{(j)} \geq \sqrt{x^2 - 1} \widetilde{V}_n^{(j)}\right) \mathbb{P}\left(|X_j| \geq \lambda\right). \tag{4.30}
\end{aligned}$$

Similarly to (4.27), we have

$$\sum_{j=1}^n \mathbb{P}\left(\widetilde{M}_n^{(j)} \geq \sqrt{x^2 - 1} \widetilde{V}_n^{(j)}\right) \mathbb{P}\left(|X_j| \geq \lambda\right) = o(1) n e^{-x^2} \tag{4.31}$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. For the first term on the right hand side of (4.30), we have

$$\begin{aligned}
\mathbb{P}\left(\widetilde{M}_n \geq x \widetilde{V}_n\right) &\geq \mathbb{P}\left(\widetilde{M}_n \geq x \widetilde{V}_n, \widetilde{V}_n^2 \leq n(1 + \varepsilon_n/x^2)\right) \\
&\geq \mathbb{P}\left(\widetilde{M}_n \geq x \sqrt{n(1 + \varepsilon_n/x^2)}\right) \\
&\quad - \mathbb{P}\left(\widetilde{M}_n \geq \sqrt{n}x, \widetilde{V}_n^2 \geq n(1 + \varepsilon_n/x^2)\right).
\end{aligned}$$

Define $\mathbf{A} = \{\widetilde{V}_n^2 \geq n(1 + \varepsilon_n/x^2)\}$. Set $Y_{k,l}(\theta_1, \theta_2) = X'_k(\theta_1 \cos(kw_l) + \theta_2 \sin(kw_l))$ for any $\theta_1, \theta_2 \in \mathbf{R}$. Let

$$\begin{aligned}\Theta_1 &= \{\theta_1 \geq 0, \theta_2 \geq 0; \theta_1^2 + \theta_2^2 = 1\}; \quad \Theta_2 = \{\theta_1 \geq 0, \theta_2 < 0; \theta_1^2 + \theta_2^2 = 1\}; \\ \Theta_3 &= \{\theta_1 < 0, \theta_2 \geq 0; \theta_1^2 + \theta_2^2 = 1\}; \quad \Theta_4 = \{\theta_1 < 0, \theta_2 < 0; \theta_1^2 + \theta_2^2 = 1\}.\end{aligned}$$

Then we have

$$\begin{aligned}\mathbb{P}(\widetilde{M}_n \geq \sqrt{n}x, \mathbf{A}) &\leq \sum_{l=1}^q \mathbb{P}\left(\sup_{(\theta_1, \theta_2) \in \Theta_1} \left| \sum_{k=1}^n Y_{k,l}(\theta_1, \theta_2) \right| \geq \sqrt{n}x, \mathbf{A}\right) \\ &\quad + \sum_{l=1}^q \mathbb{P}\left(\sup_{(\theta_1, \theta_2) \in \Theta_2} \left| \sum_{k=1}^n Y_{k,l}(\theta_1, \theta_2) \right| \geq \sqrt{n}x, \mathbf{A}\right) \\ &\quad + \sum_{l=1}^q \mathbb{P}\left(\sup_{(\theta_1, \theta_2) \in \Theta_3} \left| \sum_{k=1}^n Y_{k,l}(\theta_1, \theta_2) \right| \geq \sqrt{n}x, \mathbf{A}\right) \\ &\quad + \sum_{l=1}^q \mathbb{P}\left(\sup_{(\theta_1, \theta_2) \in \Theta_4} \left| \sum_{k=1}^n Y_{k,l}(\theta_1, \theta_2) \right| \geq \sqrt{n}x, \mathbf{A}\right).\end{aligned}\tag{4.32}$$

We only deal with the first term above, while other terms can be proved similarly. Let $\theta_{1,i} = i/n^6$ for $1 \leq i \leq n^6$ and $\theta_{2,i} = \sqrt{1 - \theta_{1,i}^2}$. We have, for $1 \leq l \leq q$,

$$\begin{aligned}\mathbb{P}\left(\sup_{(\theta_1, \theta_2) \in \Theta_1} \left| \sum_{k=1}^n Y_{k,l}(\theta_1, \theta_2) \right| \geq \sqrt{n}x, \mathbf{A}\right) &\leq \sum_{i=1}^{n^6} \mathbb{P}\left(\left| \sum_{k=1}^n Y_{k,l}(\theta_{1,i}, \theta_{2,i}) \right| \geq \sqrt{n}x - \sqrt{n}\varepsilon_n x^{-1}, \mathbf{A}\right) \\ &\quad + \sum_{i=1}^{n^6} \mathbb{P}\left(\sup_{(\theta_1, \theta_2) \in \Theta_1, \theta_{1,i-1} \leq \theta_1 \leq \theta_{1,i}} \left| \sum_{k=1}^n \left[Y_{k,l}(\theta_1, \theta_2) - Y_{k,l}(\theta_{1,i}, \theta_{2,i}) \right] \right| \geq \sqrt{n}\varepsilon_n x^{-1}\right) \\ &=: \sum_{i=1}^{n^6} J_{1,i} + \sum_{i=1}^{n^6} J_{2,i}.\end{aligned}\tag{4.33}$$

It is easy to see that $\sup_{(\theta_1, \theta_2) \in \Theta_1, \theta_{1,i-1} \leq \theta_1 \leq \theta_{1,i}} |Y_{k,l}(\theta_1, \theta_2) - Y_{k,l}(\theta_{1,i}, \theta_{2,i})| \leq n^{-1}$. Hence $J_{2,i} = 0$ for $x \in [4, \varepsilon'_n n^{1/6}]$. Letting $b = x/\sqrt{n}$ and $\tau = (\sqrt{n}/x)^{1/4}$, we have

$$\begin{aligned}J_{1,i} &\leq \mathbb{P}\left(\sum_{k=1}^n bY_{k,l}(\theta_{1,i}, \theta_{2,i}) + \tau b^2 \widetilde{V}_n^2 \geq x^2 - \varepsilon_n + \tau(x^2 + \varepsilon_n)\right) \\ &\quad + \mathbb{P}\left(\sum_{k=1}^n -bY_{k,l}(\theta_{1,i}, \theta_{2,i}) + \tau b^2 \widetilde{V}_n^2 \geq x^2 - \varepsilon_n + \tau(x^2 + \varepsilon_n)\right) \\ &\leq \mathbb{P}\left(\sum_{k=1}^n b\overline{Y}_{k,l}(\theta_{1,i}, \theta_{2,i}) + \tau b^2 [\widetilde{V}_n^2 - \mathbb{E}\widetilde{V}_n^2] \geq x^2 - \varepsilon_n + \tau\widetilde{\varepsilon}_n\right)\end{aligned}$$

$$\begin{aligned}
& +\mathbb{P}\left(\sum_{k=1}^n -b\overline{Y}_{k,l}(\theta_{1,i}, \theta_{2,i}) + \tau b^2[\widetilde{V}_n^2 - \mathbb{E}\widetilde{V}_n^2] \geq x^2 - \varepsilon_n + \tau\widetilde{\varepsilon}_n\right) \\
& =: J_{3,i},
\end{aligned} \tag{4.34}$$

where

$$\begin{aligned}
\overline{Y}_{k,l}(\theta_{1,i}) &= Y_{k,l}(\theta_{1,i}, \theta_{2,i}) - \mathbb{E}\overline{Y}_{k,l}(\theta_{1,i}, \theta_{2,i}), \\
\widetilde{\varepsilon}_n &= \varepsilon_n + \tau b^2(n - \mathbb{E}\widetilde{V}_n^2) = \varepsilon_n + o(1)x^3/\sqrt{n}.
\end{aligned}$$

Let $\eta_k = \overline{Y}_{k,l}(\theta_{1,i}, \theta_{2,i})$ and $\xi_k = X_k'^2 - \mathbb{E}X_k'^2$. Using $|e^s - 1 - s - s^2/2| \leq |s|^3 e^{s \vee 0}$, we get

$$\begin{aligned}
\mathbb{E}e^{2b\eta_k + 2\tau b^2\xi_k} &= 1 + 2\mathbb{E}(b\eta_k + \tau b^2\xi_k)^2 + O(1)\mathbb{E}|b\eta_k + \tau b^2\xi_k|^3 e^3 \\
&= 1 + 2b^2\mathbb{E}\eta_k^2 + 4\tau b^3\mathbb{E}(\eta_k\xi_k) + 2\tau^2 b^4\mathbb{E}\xi_k^2 + O(1)e^3(b^3\mathbb{E}|\eta_k|^3 + \tau^3 b^6\mathbb{E}|\xi_k|^3) \\
&= 1 + 2b^2(\mathbb{E}X_k'^2 - (\mathbb{E}X_k')^2)[\theta_{1,i}^2 \cos^2(kw_l) + \theta_{2,i}^2 \sin^2(kw_l) \\
&\quad + 2\theta_{1,i}\theta_{2,i} \cos(kw_l) \sin(kw_l)] + O(1)(1 + \tau)b^3 \\
&= 1 + 2b^2(\theta_{1,i}^2 \cos^2(kw_l) + \theta_{2,i}^2 \sin^2(kw_l) + 2\theta_{1,i}\theta_{2,i} \cos(kw_l) \sin(kw_l)) \\
&\quad + O(1)(1 + \tau)b^3,
\end{aligned}$$

for $x \in [4, \varepsilon_n' n^{1/6}]$. This, together with (4.4), implies that

$$\begin{aligned}
J_{3,i} &\leq 2 \exp\left(-2x^2 + 2\varepsilon_n - 2\tau\widetilde{\varepsilon}_n + nb^2 + O(1)(1 + \tau)x^3/\sqrt{n}\right) \\
&\leq C \exp\left(-x^2 - b^{-1/8}\right).
\end{aligned} \tag{4.35}$$

Combining (4.32)-(4.35) gives

$$\mathbb{P}\left(\widetilde{M}_n \geq \sqrt{n}x, \mathbf{A}\right) = o(1)e^{-x^2}. \tag{4.36}$$

Define

$$A_j = \left\{ \left| \sum_{k=1}^n X_k' e^{ik\omega_j} \right| \geq x\sqrt{n(1 + \varepsilon_n/x^2)} \right\}, \quad 1 \leq j \leq q.$$

We have

$$\mathbb{P}\left(\widetilde{M}_n \geq x\sqrt{n(1 + \varepsilon_n/x^2)}\right) \geq \sum_{j=1}^q \mathbb{P}(A_j) - \sum_{1 \leq i < j \leq q} \mathbb{P}(A_i A_j).$$

By Lemma 4.2 (ii), we have

$$\mathbb{P}(A_i) = (1 + o(1))e^{-x^2}, \quad \mathbb{P}(A_i A_j) = 2^{-1}(1 + o(1))e^{-2x^2},$$

uniformly in $x \in [4, \varepsilon_n' n^{1/6}]$ and $1 \leq i, j \leq q$. This shows that

$$\mathbb{P}\left(\widetilde{M}_n \geq x\sqrt{n(1 + \varepsilon_n/x^2)}\right) \geq (1 + o(1))qe^{-x^2}(1 - 2^{-1}qe^{-x^2}). \tag{4.37}$$

It follows from (4.31), (4.36) and (4.37) that

$$\mathbb{P}\left(M_n \geq xV_n\right) \geq (1 + o(1))qe^{-x^2}(1 - 2^{-1}qe^{-x^2}). \quad (4.38)$$

uniformly in $x \in [4, \varepsilon'_n n^{1/6}]$. Let $x = \sqrt{y + \log q}$. Combining (4.29) and (4.38), we have for any fixed $M > 0$,

$$\limsup_{n \rightarrow \infty} \sup_{M \leq y \leq \varepsilon'_n n^{1/3}} \left| \frac{\mathbb{P}\left(\frac{M_n^2}{V_n^2} - \log q \geq y\right)}{1 - \exp(-\exp(-y))} - 1 \right| \leq Ce^{-M}. \quad (4.39)$$

For $-\log q \leq y \leq M$, by (1.1), (4.18) and (4.19),

$$\limsup_{n \rightarrow \infty} \sup_{-\log q \leq y \leq M} \left| \frac{\mathbb{P}\left(\frac{M_n^2}{V_n^2} - \log q \geq y\right)}{1 - \exp(-\exp(-y))} - 1 \right| = 0. \quad (4.40)$$

This proves Theorem 2.3 by (4.39) and (4.40). $\square$

Proof of Lemma 3.1. This lemma follows immediately by Theorem 2.3 and

$$f_n((x + \log q)/q) = \mathbb{P}\left(\frac{\max_{1 \leq j \leq q} I_n(\omega_j)}{q^{-1} \sum_{j=1}^q I_n(\omega_j)} - \log q \geq x\right),$$

where $\{X_k\}$ are i.i.d. $N(0, 1)$ random variables. $\square$

Proof of Theorem 3.1. Let $\mathcal{C}_g = \{P_g < \theta/(3G), P_g^{true} > \theta/(2G)\}$ and define $F(x) = \exp(-\exp(-x))$. Let x_n satisfy $1 - F(x_n) = \theta/(2.5G)$. So $x_n \sim \log G$. Corollary 3.1 yields

$$\max_{1 \leq g \leq G} \left| \frac{1 - F(x_n)}{1 - F_{n,g}((x_n + \log q)/q)} - 1 \right| = o(1). \quad (4.41)$$

By (4.41) and the definition of x_n , we can see that on $\mathcal{C}_g$, it holds $P_g^{true} > \theta/(2G) > 1 - F_{n,g}((x_n + \log q)/q)$ for n large. By the monotonicity of distribution function we have $qf_g - \log q \leq x_n$. This together with Corollary 3.1 and Lemma 3.1 yields

$$\max_{1 \leq g \leq G} \left| \frac{P_g}{P_g^{true}} - 1 \right| I\{\mathcal{C}_g\} = o(1).$$

Note that on $\mathcal{H}_g \cap \mathcal{C}_g^c$ we have $P_g \geq \theta/(3G)$. We can show that $qf_g - \log q \leq y_n$, where $1 - F(y_n) = \theta/(4G)$, so $y_n \sim \log G$. In fact, by Lemma 3.1,

$$\frac{f_n(q^{-1}(y_n + \log q))}{1 - F(y_n)} - 1 = o(1),$$

and hence, $f_n(f_g) = P_g > f_n(q^{-1}(y_n + \log q))$ for n large, which implies $qf_g - \log q \leq y_n$.

It follows from Corollary 3.1 and Lemma 3.1 that

$$\max_{1 \leq g \leq G} \left| \frac{P_g}{P_g^{true}} - 1 \right| I\{\mathcal{H}_g\} I\{\mathcal{C}_g^c\} = o(1).$$

The theorem is proved. $\square$

References

- [1] Ahdesmäki, M., Lähdesmäki, H., Pearson, R., Huttunen, H. and Yli-Harja, O. (2005). Robust detection of periodic time series measured from biological systems. *BMC Bioinformatics*, **6**: 117 1-18.
- [2] Amosova, N.N. (1982). Probabilities of moderate deviations. *J. Math. Sci.* **20**: 2123-2130.
- [3] An, H. Z., Chen, Z. G. and Hannan, E. J. (1983). The maximum of the periodogram. *J. Multivariate Anal.* **13**: 383-400.
- [4] Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B* **57**: 289-300.
- [5] Chen, J. (2005). Identification of significant periodic genes in microarray gene expression data. *BMC Bioinformatics*, **6**: 286 1-12.
- [6] Davis, R. A. and Mikosch, T. (1999). The maximum of the periodogram of a non-Gaussian sequence. *Ann. Probab.* **27**: 522-536.
- [7] de la Peña, V.H., Lai, T.L. and Shao, Q.M. (2009). Self-normalized processes: limit theory and statistical applications. Springer, New York.
- [8] Fan, J.Q., Hall, P and Yao, Q. (2007). To how many simultaneous hypothesis tests can normal, student's t or bootstrap calibration be applied? *J. Am. Stat. Assoc.* **102**:1282-1288.
- [9] Fay, G. and Soulier, P. (2001). The periodogram of an i.i.d. sequence. *Stoch. Proc. Appl.* **92**: 315-343.
- [10] Fisher, R. A. (1929). Tests of significance in harmonic analysis. *Proc. Roy. Statist. Soc. Ser. A* **125**: 54-59.
- [11] Glynn, E., Chen, J. and Mushegian, A. (2006). Detecting periodic patterns in unevenly spaced gene expression time series using Lomb-Scargle periodograms. *Bioinformatics*, **22(3)**: 310-316.
- [12] Hu, Z., Shao, Q.M. and Wang, Q. (2009). Cramér-type moderate deviations for the maximum of self-normalized sums. *Elect. J. Probab.* **14**: 1181-1197.
- [13] Jing, B.Y., Shao, Q.M. and Wang, Q. (2003). Self-normalized Cramér-type large deviations for independent random variables. *Ann. Probab.* **31**: 2167-2215.
- [14] Korosok, M.R. and Ma, S. (2007), Marginal asymptotics for the large p , small n paradigm: with applications to microarray data. *Ann. Statist.* **35**: 1456-1486.
- [15] Ledoux, M. and Talagrand, M. (1991). Probability in Banach spaces. Springer, Berlin.

- [16] Lin, Z.Y. and Liu, W.D. (2009a). On maxima of periodograms of stationary processes. *Ann. Statist.* **37**: 2676-2695.
- [17] Lin, Z.Y. and Liu, W.D. (2009b). Supplementary material for "On maxima of periodograms of stationary processes." Available at <http://xxx.arxiv.org/abs/0801.1357>.
- [18] Michel, R. (1976). Nonuniform central limit bounds with application to probabilities of deviations. *Ann. Probab.* **4**: 102-106.
- [19] Mikosch, T., Resnick, S., and Samorodnitsky, G. (2000). The maximum of the periodogram for a heavy-tailed sequence. *Ann. Probab.* **28**: 885-908.
- [20] Petrov, V.V. (2002). On probabilities of moderate deviations. *J. Math. Sci.* **109**: 2189-2191.
- [21] Shao, Q. M. (1997). Self-normalized large deviations. *Ann. Probab.* **25**: 285-328.
- [22] Shao, Q. M. (1999). A Cramér type large deviation for Student's t statistic. *J. Theoret. Probab.* **12**: 385-398.
- [23] Shao, X. and Wu, W. B. (2007). Asymptotic spectral theory for nonlinear time series. *Ann. Statist.* **35 (4)**: 1773-1801.
- [24] Wang, Q. (2005). Limit theorems for self-normalized large deviations. *Elect. J. Probab.* **10**: 1260-1285.
- [25] Wichert, S., Fokianos, K. and Strimmer, K. (2004). Identifying periodically expressed transcripts in microarray time series data. *Bioinformatics*, **20(1)**: 5-20.
- [26] Wu, W. B. and Zhao, Z. (2008). Moderate deviations for stationary processes. *Statist. Sinica*. **18**: 769-782.